# 2023 MHQ Summary

Client Name





## Screening Tools Used

The Mental Health Questionnaire was created by combining four gold standard mental health assessments, as outlined below.

PHQ-9

An instrument for screening, diagnosing, monitoring, and measuring the severity of depression according to DSM-IV depression diagnostic criteria.

GAD-7

A screening tool that uses some of the DSM-V criteria for GAD (General Anxiety Disorder) to identify probable cases of GAD along with measuring anxiety symptom severity.

PC-PTSD-5

A 5-item screen designed to identify individuals with probable PTSD.


CAGE


CAGE is an internationally used assessment instrument for identifying alcoholics.




#### Demographics

22 respondents9% of employee populationNOT a representative sample sizeN/A = "I prefer not to answer"










### PHQ-9



- **50%** of respondents experience at least 1 symptom of depression
- 2/3 respondents experiencing mild depression have symptoms at the higher end of the range, i.e. closer to moderate depression
- 2 respondents experience moderately severe depression

**67%** 

of Dearborn respondents experience symptoms of depression

ALL


age ranges are represented among those who screened positively for some level of depression

100%

of the women respondents screened positively for some level of depression



### GAD-7



- **59%** of respondents experience at least 1 symptom of anxiety
- of those who experience some level of anxiety, over half experience minimal symptoms
- 3 respondents experience severe anxiety
- of the 3 who experience severe anxiety, **2** also experience moderately severe depression


of respondents who screened positively for some level of depression ALSO screened positively for some level of anxiety

of the men respondents screened positively for some level of anxiety

of the women respondents screened positively for some level of anxiety




### PC-PTSD-5



- 22% of respondents indicated that they had experienced a traumatic event
- 2/5 were below the threshold
- **3** (14% of respondents) optimally sensitive to probable PTSD
  - o all 3 experience some level of depression
  - of the 3, 2 also experience severe anxiety
  - of the 3, 1 also experiences severe anxiety and moderately severe depression



### CAGE



- 1 clinically significant response
  - this individual also experiences mild anxiety and minimal depression
- Have you ever felt you should **C**ut down on your drinking?
  - 3 respondents answered "Yes"
- A Have people Annoyed you by criticizing your drinking?
  - 1 respondent answered "Yes"
- Have you ever felt bad or **G**uilty about your drinking?
  - 1 respondent answered "Yes"
- Have you ever had a drink first thing in the morning to steady your nerves or to get rid of a hangover (**E**ye opener)?

